

The neurobiological mechanisms of physical exercise in methamphetamine addiction

António Pedro Delgado Morais¹ | Inês Roque Pita¹ | Carlos Alberto Fontes-Ribeiro^{1,2} | Frederico Costa Pereira^{1,2} 

¹Institute of Pharmacology and Experimental Therapeutics/IBILI, Faculty of Medicine, University of Coimbra, Coimbra, Portugal

²CNC.IBILI—University of Coimbra, Coimbra, Portugal

Correspondence

Frederico Costa Pereira, Lab. Pharmacology and Experimental Therapeutics/IBILI, Faculty of Medicine, University of Coimbra, Coimbra, Portugal.

Email: fredcp@ci.uc.pt

Funding information

Fundação para a Ciência e Tecnologia (FCT, Portugal), Grant/Award Number: UID/NEU/04539/2013; COMPETE-FEDER, Grant/Award Number: POCI-01-0145-FEDER-007400; Centro 2020 Regional Operational Programmes, Grant/Award Number: CENTRO-01-0145-FEDER-000012 and CENTRO-01-0145-FEDER-000008

Abstract

Methamphetamine (METH) is the primary drug within amphetamine-type stimulants which are the second most abused group of drugs worldwide. There is no pharmacological treatment addressed specifically to METH addiction, and behavioral therapy is shadowed by poor long-term recovery and relapse. Therefore, novel approaches to manage METH addiction are an urgent need. This review aims to describe the current state of physical exercise use on methamphetamine addiction management. The following searching terms in PubMed were used: ("physical exercise" OR "exercise") AND "methamphetamine." Relevant references from key publications and gray literature were also reviewed to identify additional citations for inclusion. Original investigation regarding physical exercise and methamphetamine addiction (clinical data) or neurobiological mechanisms of physical exercise in animal models of methamphetamine administration (preclinical data) was included. Overall, METH users demonstrated improvements, including better fitness and emotional measures, lower relapse rates, and sustained abstinence when compared to nonexercised individuals. The neurobiological mechanisms of physical exercise in METH users seem to reflect an interplay of several agents, including neurochemicals, oxidative stress, neurogenesis, gliogenesis, and blood-brain barrier as disclosed by preclinical data. Exercise-based interventions alone or as a conjoint therapy may be a useful tool for managing METH addiction.

KEYWORDS

amphetamine-type stimulants, methamphetamine, methamphetamine addiction, physical exercise

1 | INTRODUCTION

The most recent report from United Nations Office on Drugs and Crime indicates that amphetamine-type stimulants (ATS) are the second most abused group of drugs worldwide.¹ After 3 years of relative stability, ATS seizures reached a new peak of more than 170 tons in 2014.¹ For the past few years, methamphetamine (METH) seizures have accounted for the largest share of global ATS seizures annually (Figure 1), but, although METH is a feature of ATS markets worldwide, it is particularly dominant in East and South-East Asia and North

America.¹ It is estimated that 29 million people suffer from drug use disorders, but only 1 in 6 is in treatment.¹ Importantly, the number of people requiring treatment for ATS use is increasing.¹ METH use, in particular, accounts for a large share of people receiving drug treatment in large parts of East and South-East Asia.¹ METH users seeking treatment is also increasing in certain parts of the United States, as well as deaths related to METH misuse²⁻⁴ (Figure 2). Therefore, METH use is a serious worldwide public health problem with major psychiatric and medical consequences, including psychosis, dependence, overdose/death, cognitive, socioeconomic, and legal consequences.³

The first two authors contributed equally to this work.

Global seizures

Change from previous year




FIGURE 1 Global seizures of ATS including methamphetamine, amphetamine, and "ecstasy" in 2014. This figure was adapted from United Nations Office on Drugs and Crime (UNODC), 2016—World Drug Report 2016




FIGURE 2 Drugs involved in U.S. overdose deaths, 2000-2016. Figure adapted from <https://www.drugabuse.gov/related-topics/trends-statistics/overdose-death-rates>

However, no medication has been approved by the regulatory authorities for the treatment of METH addiction. Finding effective treatment for this clinical problem is certainly an urgent unmet need. The aim of this review was to summarize the evidence of the employment of physical exercise in METH addiction management.

2 | METHODOLOGY

2.1 | Search, information source, and selection studies

A search in the online database PubMed was carried out updated until April 5, 2017, using the following search terms ("methamphetamine") AND (("physical exercise") OR ("exercise")). A total of 101 full-text articles were obtained on which inclusion and exclusion criteria were applied. Publications were included if (i) controlled clinical studies evaluating exercise as a therapeutic intervention for METH addiction were reported or if (ii) preclinical studies using METH-administration

models subjected to physical exercise were reported and if (iii) papers were written in English. Publications were excluded if (i) the study design was a review or (ii) METH was used only once or (iii) physical activity was used as a readout of METH neurotoxicity. Twenty studies were included after applying the inclusion and exclusion criteria. Gray literature retrieved from U.S. Department of Justice Drug Enforcement Administration and from United Nations was also considered. Finally, relevant references from key publications obtained from the PubMed search were also included.

3 | METH CLINICAL FEATURES

3.1 | METH pharmacokinetics

METH can be smoked, injected, ingested, dissolved sublingually, or solubilized and when consumed as a liquid, inserted rectally or into the urethra.³ Smoking (inhaling the fumes after heating the hydrochloride form) is the most common route of administration and has the highest bioavailability following injection (dissolving the hydrochloride form in water prior to injection).⁵⁻⁷ When smoked or injected intravenously, METH rapidly reaches the cerebral circulation (6-15 seconds), and a rapid onset of euphoria ensues, which typically lasts for several minutes.⁸ Therefore, these two routes carry higher potential for acute toxicity, addiction and overdose.⁸ Moreover, the risk of infectious diseases, such as human immunodeficiency virus (HIV) and viral hepatitis, through unsafe injecting practices, is noteworthy.⁹ In opposition, when "snorted" or ingested in powder form, METH takes approximately 5 and 20 minutes, respectively, to reach a peak of euphoric state.¹⁰ However, this effect can last from 8 to 12 hours.

The metabolism of METH occurs largely in the liver via phase I reactions catalyzed by cytochrome CYP2D6, which produces amphetamine, 4-hydroxymethamphetamine and norephedrine, among other metabolites unlikely to influence the clinical spectrum.¹¹ Approximately 70% of the drug is then excreted by the kidneys, mainly as nonmetabolized METH (30%-50%), followed by up to 15% as 4-hydroxymethamphetamine and 10% as amphetamine.^{7,12}

3.2 | METH toxicodynamics

METH is a cationic lipophilic molecule, being the most potent ATS drug.¹³ METH primarily causes the release of the monoamines dopamine (DA), serotonin (5-HT), and norepinephrine (NA). NA is released most efficiently, followed by DA and then 5-HT.^{14,15} A wealth of studies has focused on METH impact on DA release, due to this monoamine's crucial role in reward and reinforcement processes.^{11,16} This psychostimulant triggers an aberrant release of DA from the presynaptic terminal into the synaptic cleft as the outcome of the complex interplay of the primary mechanisms of action of METH: the redistribution of DA from synaptic vesicles to the cytosol through the inhibition of vesicular monoamine transporter-2 (VMAT-2) and the reverse transport of DA via DA transporter (DAT).^{11,14} METH also competitively inhibits DA uptake and monoamine oxidase.^{14,17}

When taken at low-to-moderate doses (5–30 mg), METH induces short-term symptoms that are related with a sympathetic response by the autonomous nervous system (tachycardia, tachypnea, hypertension, pupil dilation, hyperthermia, reduced fatigue) and include a state of euphoria and social ease of relating with other people, increased attention, behavioral disinhibition, reduced appetite, and sense of increased energy, sex drive, and self-confidence.^{3,8} Repeated METH abuse results in central nervous system (CNS) disturbances that can range from insomnia, aggression, and mood/anxiety symptoms to severe neuropsychiatric changes.¹⁸ Chronic METH users generally present a decline in cognition associated with impaired episodic memory, executive functions, complex information processing speed, and psychomotor functions¹⁹ that can persist during early²⁰ and prolonged abstinence.²¹ METH users may also present psychosis, comparable to acute episodes of schizophrenia, in which users report delusions, hallucinations, and odd speech.^{3,8}

There has been much effort put in elucidating the structural, neurochemical, and metabolic substrate for this well-characterized METH neurotoxicity. Functional neuroimaging studies have documented several alterations in brain activation patterns induced by METH.²² For example, METH users who showed impaired attention and impaired cognitive control exhibited abnormalities in cingulate gyrus and insula. Additionally, high-resolution MRI and surface-based computational image analyses generated cortical maps that revealed severe gray matter deficits in the cingulate, limbic, and paralimbic cortices of MA abusers, smaller hippocampal volumes than control subjects, and significant white matter hypertrophy.²³ Tobias et al²⁴ further showed METH abuse produces microstructural abnormalities in white matter underlying and interconnecting prefrontal cortices and hippocampal formation. Furthermore, decreased frontal activation associated with impaired decision making and cognitive control was also disclosed in METH users.^{22,25} Maladaptive decision making by METH users may reflect circuit-level dysfunction, underlying deficits in task-based activation: Increased resting-state connectivity within the mesocortico-limbic system coupled with reduced prefrontal cortical connectivity may create a bias toward reward-driven behavior over cognitive control in methamphetamine users.²⁶

Additionally, METH abusers have abnormalities in brain regions implicated in mood disorders. In fact, self-reports of depressive symptoms covaried positively with relative glucose metabolism in limbic regions (eg, perigenual anterior cingulate gyrus and amygdala) and ratings of state and trait anxiety covaried negatively with relative activity in the anterior cingulate cortex, orbitofrontal cortex, and left insula, and positively with amygdala activity.²⁷ Functional magnetic resonance imaging (fMRI) studies disclosed cortical abnormalities that could underlie the socially inappropriate behaviors including violence and aggression often shown by individuals who abuse METH.²⁸ London et al²⁹ reviewed clinical studies indicating chronic methamphetamine users often display several signs of corticostriatal dysfunction, including abnormal gray and white matter integrity, monoamine neurotransmitter system deficiencies, neuroinflammation, poor neuronal integrity, and aberrant patterns of brain connectivity and function, both when engaged in cognitive tasks and at rest.

Regarding neurochemical data, clinical studies indicated the presence of a marked monoaminergic dysfunction as reflected by a down-regulation of VMAT-2, 5-HT transporter (SERT), DA D2 receptor and DAT in the striatum (nucleus accumbens, caudate and putamen),^{21,30,31} and orbitofrontal and dorsolateral prefrontal cortex³² of METH abusers. It was shown that low striatal D(2)/D(3) receptor availability may mediate impulsive temperament and thereby influence addiction in METH-dependent subjects.³³ Moreover, this low striatal DA receptor availability was linked to caloric intake during abstinence from chronic methamphetamine abuse.³⁴ Postmortem studies also provided evidence of low levels of DA, tyrosine hydroxylase (TH), and DAT in the striatum^{35–37} and SERT in orbitofrontal and occipital cortices.³⁸

Oxidative stress is another culprit of METH-induced neurotoxicity. In fact, studies in humans demonstrated that prolonged use of METH exerted oxidative stress and enhanced lipid peroxidation, as METH abusers had significantly higher levels of plasmatic malondialdehyde (MDA, a lipid peroxidation product), when compared to healthy controls.³⁹ Moreover, METH abusers have persistently higher systemic oxidative stress throughout early abstinence as levered by higher plasmatic levels of MDA, which were accompanied by lower activity of superoxide dismutase (SOD), and higher activity of catalase and levels of glutathione relatively to healthy controls.

New and emergent aspects of METH toxicity involve blood-brain barrier (BBB), neurogenesis, and gliogenesis dysfunction. The majority of evidence regarding BBB disruption is provided by preclinical studies.^{40–43} There is scarce human literature corroborating these preclinical evidences. For example, Ago et al⁴⁴ reported that a man, who intravenously injected METH, showed brain edema. Additionally, preclinical data suggest that METH can also affect neurogenesis and gliogenesis. For example, Mandyam et al⁴⁵ showed that self-administration with daily short or extended access to METH (which mimics a human recreational use) generated a dependence-like syndrome and decreased gliogenesis in medial prefrontal cortex from rats. Mice that were subjected to METH administration for 5 days with an escalating dose regimen showed a reduction in the number of differentiating neural cells in the hippocampal dentate gyrus (DG).⁴⁶ In rats self-administering METH for 28 days, the number of proliferating neural progenitor cells

in the hippocampal DGs was significantly increased by METH exposure for 1 hour twice weekly, not changed by daily self-administration for 1 hour, and significantly decreased by daily exposure for 6 hours.⁴⁷ Importantly, a robust rebound in neurogenesis was associated with enhanced propensity for reinstatement during METH abstinence in METH self-administered rats.⁴⁷ Moreover, forced abstinence from higher preferred levels of METH intake enhanced neurogenesis and neuronal activation of granule cell neurons (GCNs) in the DG and produced compulsive-like drug reinstatement in rats.⁴⁸ Furthermore, systemic treatment with the drug isoxazole-9 (a synthetic small molecule known to modulate neurogenesis in the adult rodent brain) during abstinence blocked compulsive-like context-driven methamphetamine reinstatement.⁴⁸ These results suggest that adult neurogenesis during abstinence play a role in compulsive-like METH reinstatement.

METH use produces also serious complications in multiple organs, beyond the brain.^{49,50} In fact, hyperthermia, hypertension, cardiac arrhythmia, seizures, cerebral hemorrhage, ischemic infarct, renal failure, liver damage, rhabdomyolysis, and wakefulness to the point of collapse and temporary blindness, coma, or death may ensue when METH overdose is in place.⁵¹⁻⁵⁵

Similarly to other drugs of abuse, sudden cessation of METH consumption causes a withdrawal syndrome presenting as anhedonia, irritable or aggressive mood, anxiety, craving, sleep disturbance, diminished cognitive functions, and musculoskeletal pain, among other signs and symptoms, with a prevalence of depressive symptoms.^{3,56} This withdrawal syndrome may be a consequence of overall METH neurotoxicity.¹¹ The neuropsychological consequences of this intricate and complex METH neurotoxicity render management of METH addiction a challenging task.

4 | BIOLOGICAL SEX AND METH ADDICTION

There is scarce information about gender differences in (METH)-dependent users. However, women and men reported regular METH use at a similar age, but, women reported experiencing problems as a result of METH use at an earlier age than men, suggesting a more rapid progression from initial drug use to chronic METH use, and then to problem use.^{57,58} A recent epidemiological survey suggested that the proportions of female and male admissions reporting methamphetamine/amphetamines as their primary substance of abuse were similar across all age groups with the exception of those aged 18-24. Specifically, among admissions aged 18-24, 8.9 percent of female admissions reported primary methamphetamine/amphetamine abuse compared with 3.7% of male admissions.⁵⁹ Regarding reasons for using METH, women reported "increased energy" and "losing weight", whereas men reported using METH to "complete more work" or to "try something new".⁶⁰ A recent study disclosed that METH-dependent women had greater psychological burden, reported more use of an emotional-coping strategy, and had greater childhood emotional and sexual trauma.⁶¹ Kogachi et al⁶² investigated the relationships between impulsivity, brain structures, and possible sex-specific

differences between METH users and non-drug-using controls. There was no difference in impulsivity between the male and female METH users. However, the female METH users who started using METH at an earlier age had higher impulsivity scores. Contrary to the male METH users, female METH users had smaller and thinner frontal cortices. Moreover, cortical changes in both sexes were associated with greater cognitive impulsivity. These data suggest that sex may modulate the effects of METH on brain morphometry. Overall, the need for gender-tailored METH use treatment programs as suggested by Simpson et al⁶¹ is seemingly warranted.

5 | CURRENT MANAGEMENT AND TREATMENT OF METH ADDICTION

Treatment of METH addiction is a major problem, with elevated overall expenditures due to the requirement of specialized personnel and infrastructures, along with the constant monitoring and follow-up of these patients.⁶³ Currently, there are limited therapeutic options, with low efficacy and high relapsing rates. Treatment can be directed to acute syndromes or chronic METH abuse. The purpose of this review is to focus on the chronic form of METH addiction.

Methamphetamine user patterns need to be taken into account, when managing METH abusers. Low-, moderate-, and high-use patterns relate with treatment trajectories, as greater treatment participation was achieved when METH use frequency was lower.⁶⁴ In addition, individuals who initiated METH consumption at a later age tended to look for treatment at an earlier stage and are less inclined to inject themselves than those who started earlier. Besides, polydrug abuse, often seen on early set use, contributes to low-treatment compliance. An early intervention in all patterns of abuse is essential to decrease METH intake. Gender may also influence treatment outcomes. A follow-up study noticed that, although women were faced with more challenges during treatment, such as unemployment or psychiatric symptoms, compared to men, they were capable of a greater improvement in several areas, including family relationships and medical problems.⁵⁷

In addition, METH withdrawal syndrome is another determinant factor for the treatment process. In fact, the neuropsychiatric impairments (eg, cognitive deficits, depression, and anxiety) that persist following abstinence are associated with poorer treatment outcomes, as follows: increased relapse rates, lower treatment retention rates, and reduced daily functioning.⁶⁵ Brecht and Herbeck⁶⁶ evaluated the relapsing rates for periods averaging 5 years postdischarge from treatment for METH use. These authors found that, in a sample of 350 subjects, only 13% remained abstinent in the whole 5-year period, and 61% relapsed within the first-year posttreatment, with half the sample relapsing within 6 months. Treatment modality was residential for 62% and outpatient for 38% of METH abusers. Relapsing rates decreased as abstinence period increased. Having experienced serious METH-related psychiatric/behavioral problems and longer duration of treatment worked as protective factors against relapse, while risk factors included having a drug addicted parent and involvement

in METH trade. Initial abstinence should be addressed with supportive measures, such as healthy eating, resting, and exercising, with this one adding a promising new approach.⁶⁷ These measures may aid pharmacotherapy, which is mostly unsuccessful, and psychosocial/behavioral treatment, the main, yet flawed, strategy regarding METH dependence.³

5.1 | Pharmacotherapy

So far, there is no specific pharmacological treatment for METH addiction.^{3,68} Guidelines on this subject regard former and current experience with other stimulant's dependence and they lack proven efficacy studies.⁸

Pharmacological approaches evaluated for METH addiction may interfere with the dopaminergic reward pathways, attenuate negative reinforcing effects of withdrawal, or improve psychiatric symptoms that impair chances of remaining abstinent.⁶⁹ This leads to several types of medications that were tested in human laboratory paradigms and in Phase II clinical trials with different outcomes as follows: dexamphetamine, methylphenidate, modafinil (DA indirectly-acting agonists), bupropion, mirtazapine (antidepressants), aripiprazole (antipsychotics), varenicline (nicotinic receptor partial agonist), rivastigmine (cholinesterase inhibitor), perindopril (angiotensin-converting enzyme inhibitor IECA), ibudilast (a phosphodiesterase inhibitor; glial modulator), baclofen, gabapentin (GABAergic agents), topiramate (GABAergic/glutamatergic agent), naltrexone (opioid antagonist), n-acetylcysteine (NAC, antioxidant), ondansetron (5-HT₃ receptor antagonist), passive immunization with monoclonal antibodies against METH (ie, vaccines).^{69–86} Worley et al⁷¹ highlighted that mirtazapine, bupropion, and methylphenidate have shown some beneficial effect in phase II clinical trials, whereas mirtazapine and bupropion increased METH abstinence, and methylphenidate reduced METH craving and use and improved depressive symptoms. However, the overall efficacy is low. We argue that the physical exercise could augment the efficacy of pharmacotherapy, thus providing a broader and stronger effect in the METH-dependent population. In fact, it was already shown that the combination of PE and sertraline could improve the management of late-life depression, especially when customized for individuals with specific clinical features.⁷² The combination of physical exercise and pharmacotherapy in the context of METH addiction needs to be tested in clinical trials.

5.2 | Psychosocial Therapy

Cognitive and behavioral treatments consist essentially in cognitive-behavioral therapy (CBT) and contingency management (CM) and generated good clinical outcomes in METH users.⁸⁷ CBT comprises several structure sessions led by a therapist which intends to raise self-awareness to negative actions or particular situations. Although CBT decreases stimulant use, this decrease is not as significant as in CM. The matrix model, a derivative of CBT, was created in the early 1980s to treat cocaine dependence and has suffered several changes since then.⁸⁸ Their overall goals consist in the following: interrupting

drug abuse, understanding the issues behind relapse and addiction, providing support and education to family members, familiarizing with self-help programs, and monitoring the follow-up status by, for example, collecting urine samples.⁶³ The program lasts for a preestablished period of 16 weeks and it has, generally, successful outcomes.

Contingency management was originally used in opiate abusers, and it works by strengthening positive behaviors by rewarding the patient, thus leading to sustained abstinence. Rewards include vouchers that can be exchanged for food or any other item or service and even cash incentives. When compared to a counseling-only strategy, CM yielded less positive urine samples and longer periods of abstinence.⁶⁷

Behavioral therapies can be part of a drug court treatment, with the verdict of remain "clean" for at least 1 year and according to a number of norms. Indeed, users who attended drug court treatment fulfilled greater rates of abstinence, retention, and completion.⁸⁹ However, Courtney and Ray³ suggested that caveats must be considered when interpreting conclusions regarding positive outcomes from these psychosocial approaches. For example, the durability of treatment effects (especially with respect to CM programs) was highlighted. Additionally, the effectiveness of psychosocial interventions may be shadowed by poor rates of treatment induction and retention.⁹⁰ Furthermore, METH-related cognitive deficits related to inhibitory control may potentially hamper the efficacy of heavily cognitive-based treatments.^{3,91}

6 | CLINICAL EVIDENCE FOR PHYSICAL EXERCISE ON METH ADDICTION

Physical exercise has been proposed as a potential treatment for METH addiction. Indeed, exercise was shown to attenuate symptoms of depression and anxiety, prevent addictive behaviors, and improve sleep and cognitive impairments, features often observed in current and former METH users.^{92,93} Clinical positive outcomes were observed for several substances, both legal, as tobacco⁹⁴ and alcohol,⁹⁵ and illegal, as cannabis⁹⁶ and heroin.⁹⁷

Newly abstinent METH users submitted to an 8-week endurance training (3 days/week for 3 weeks of jogging and/or walking on a treadmill for 30 minutes at an intensity based on heart rate plus 5 weeks of increasing intensity) followed by a resistance program (progressive, circuit-type program using selectorized machines, and/or dumbbell training that included all the major muscle groups of the upper and lower body) showed substantial improvements in aerobic exercise performance, muscle strength and endurance, body composition, and increased heart rate variability as an index of a healthy autonomic nervous system, which together could enhance recovery from drug dependency.^{98,99} However, these authors did not assess endpoints directly associated with METH addiction. Robertson et al¹⁰⁰ aimed to evaluate whether adding an exercise training program to an inpatient behavioral intervention for METH addiction reversed deficits in striatal D₂-type receptors. Participants remained abstinent from drugs and received behavioral therapy for their addiction plus 1 h of supervised exercise training, 3 days/week for 8 weeks as implemented by Dolezal et al^{98,99} Control participants received equal-time

TABLE 1 Clinical evidence of physical exercise use on methamphetamine addiction management

Reference	Sample	Type of Exercise	Duration of exercise	Outcomes
Dolezal et al ⁹⁸	29 METH-dependent individuals finished the proposed program, resulting in a 74% adherence rate; 15 elements in the exercise group and 14 in a health education group without training (sedentary) (n = 39 individuals)	<i>Aerobic training</i> (30 min): first 3 wks jogging and/or walking on treadmill during 30 min, at intensity based on heart rate (HR); the subsequent 5 wks had increasing intensity; <i>Resistance training</i> (30 min): progressive, circuit-type, resistance training that included all the major muscle groups of the upper and lower body	3 d/wk; 8 wks	Aerobic capacity and endurance: improvement of VO2 max (\uparrow 21%) as well as muscle strength and endurance in the exercise group; Body composition and anthropometry: reduction in percent relative body fat (\downarrow 15%) and reduction in fat weight (\downarrow 18%) in the exercise group.
Dolezal et al ⁹⁹	28 METH-dependent individuals under residential treatment were divided into two subgroups of 14 elements: exercise and equal-attention health education program without training (sedentary); these were compared to 22 aged-matched, drug-free, sedentary controls (n = 50, all males)	<i>Aerobic training</i> (30 min): first 3 wks jogging and/or walking on treadmill during 30 min, at intensity based on HR; the subsequent 5 wks had increasing intensity; <i>Resistance training</i> (30 min): progressive, circuit-type, resistance training that included all the major muscle groups of the upper and lower body	3 d/wk; 8 wks	Exercise markedly increased HRV (HRV was reduced in recently abstinent users when compared with sedentary drug-free controls); Exercise improved gain in aerobic capacity (VO2 max; 24%); Exercise increased muscle strength and endurance for upper (51% and 90%, respectively) and lower body (40% and 112%, respectively) (resistance training); Body composition and anthropometry: reductions in body mass ($-$ 3%), percent relative body fat ($-$ 14%), and body mass index ($-$ 4%) in the exercise group.
Rawson et al ⁹³	138 METH-dependent individuals under residential treatment were randomly assigned to the exercise group (67 users) or to the Educational group (sedentary; 71 users)	<i>Aerobic training</i> : 5 min of warm-up, 30 min of aerobic activity on a treadmill; <i>Resistance training</i> : 15 min weight lifting in major muscle groups, and 5 min of cooldown and stretching; 3 d/wk	3 d/wk, 8 wks	8-week follow-up postdischarge: 1 and 2 patients were lost to follow-up from the sedentary and the exercise group, respectively; Physical exercise significantly reduced depression and anxiety symptom scores at study discharge (according to Beck Depression Inventory)
Rawson et al ¹⁰¹	138 METH-dependent individuals under residential treatment were randomly assigned to the exercise group (71 users) or the educational group (sedentary) (67 users)	<i>Aerobic training</i> : 5 min of warm-up, 30 min of aerobic activity on a treadmill; <i>Resistance training</i> : 15 min weightlifting in major muscle groups, and 5 min of cooldown and stretching; 3 d/wk	3 d/wk, 8 wks	8-week follow-up postdischarge: 1 and 2 patients were lost to follow-up from the sedentary and the exercise group, respectively; Physical exercise (followed by no additional encouragement or support for continued exercise) decreased METH use among lower severity METH users at 1, 3, and 6 mo posttreatment. This benefit was sustained for 6 mo.
Robertson et al ¹⁰⁰	METH-dependent individuals were randomized to a group that received 1 h supervised exercise training (exercised; n = 10) or one that received equal-time health education training (sedentary; n = 9)	<i>Aerobic training</i> (30 min): first 3 wks jogging and/or walking on treadmill during 30 min, at intensity based on HR; the subsequent 5 wks had increasing intensity; <i>Resistance training</i> (30 min): progressive, circuit-type, resistance training that included all the major muscle groups of the upper and lower body	3 d/wk; 8 wks	Exercised patients displayed a significant increase in striatal D2/D3 receptor availability compared to the sedentary group; There were no changes in D2/D3 receptor availability in extrastriatal regions in either group.

(Continues)

TABLE 1 (Continued)

Reference	Sample	Type of Exercise	Duration of exercise	Outcomes
Wang et al ¹⁰³	Participants were randomly divided into the four treatment groups: the three intensities of exercise treatments (light, moderate, and high) and a reading control	Aerobic exercise: 5-min warm-up; 20-min exercise using a bicycle ergometer at 50 rpm; 5-min cooldown; Participants were instructed to cycle while keeping their HR at one of three desired exercise intensities: within the range of 40%-50%, 65%-75%, or 85%-95% of their maximum HR	Two bouts of acute aerobic exercise 1-week apart	Acute moderate-intensity exercise may be associated with more positive effects related with METH-associated craving and inhibitory control in METH-dependent individuals (behavioral and neuroelectric measures).
Wang et al ¹⁰⁵	Sixty-two people with METH dependencies were recruited through the Drug Rehabilitation Bureau and were assigned to either an aerobic exercise or attentional control group; 50 participants completed the trial.	Aerobic exercise: 5-min warm-up; 30-min sessions of moderate-intensity exercise (ie, cycling, jogging, or jump rope); 5-min cooldown; The exercise training program began at an intensity of 65%-70% of the HR max for each METH-dependent individual. After the second week, the intensity was gradually increased to 70%-75% of HR max, based on the participant's response.	3 d/wk; 12 wks	Moderate-intensity aerobic exercise training attenuated METH-associated cravings and improved inhibitory control in METH-dependent individuals (behavioral and neuroelectric measures).

health education training. D2/D3 receptor availability [measured as nondisplaceable binding potential (BPND)] was monitored by positron emission tomography (PET) using [(18F)]fallypride. Structured exercise training was able to increase striatal D2/D3 receptor availability in comparison to equal time of health education training. These authors suggested further evaluation of physical exercise as an adjunctive treatment for stimulant dependence is warranted.

Other studies also subjected METH-dependent adults to a 8-week period moderate exercise program [5 minutes of warm-up, 30 minutes of aerobic activity on a treadmill (endurance), 15 minutes of resistance training with weightlifting in major muscle groups, and 5 minutes of cooldown and stretching; 3 days/week]. Treadmill speed and resistance weight were progressively increased. The exercised participants showed less depression and anxiety symptoms and returned less to METH use postdischarge, compared to health education participants.^{93,101,102} Health education sessions comprised wellness multimedia classes, including nutrition, sleep hygiene, time management, or health screening recommendations. Haglund et al¹⁰² further stressed that participants who attended the greatest number of exercise sessions derived the greatest benefit. Furthermore, exercise appeared to be particularly beneficial to individuals who suffered from severe medical, psychiatric, and addictive disorders.¹⁰²

While physical exercise has shown efficacy in correcting emotional deficits in METH users, METH craving is also a tempting target. Notably, Wang et al¹⁰³ expanded Wang et al¹⁰⁴ observations and further showed that acute aerobic exercise (5 minutes warm-up, a 20-minute main exercise on a stationary cycle ergometer at 50 RPMs, and a 5-minute cooldown) may provide benefits for METH-associated cravings and inhibitory control (which is referred as the ability to control inadequate behavior), as revealed by behavioral and neuroelectric measures. Moderate-intensity exercise (heart rate within 65%-75% of the participant's estimated maximum value) may be associated with more positive effects when compared with estimated maximum value and may be associated with more positive effects when compared with light and vigorous intensity. Wang et al¹⁰⁵ extended these observations by demonstrating that aerobic exercise training [a 12-week program: 5-minute warm-up; 30-minute sessions of moderate-intensity exercise (ie, cycling, jogging, or jump rope; heart rate within 65%-75% of the participant's estimated maximum value); 5-minute cooldown] may be also efficacious for METH-associated cravings and inhibitory control among METH-dependent individuals. These authors argued that these evidences may contribute to the development of specific exercise prescriptions for special populations. Table 1 summarizes the clinical studies showing evidence of physical exercise usefulness in managing METH addiction.

Regarding the growing number of studies for the last few years, we argue that exercise-based interventions for managing METH abuse are indeed highly auspicious. Many forms of exercise (eg, running and swimming) may be conducted independently, either at home or outdoors and have the potential to be cost-effective, flexible, and accessible.¹⁰⁶

However, three cautionary notes should be added. Firstly, there could be various challenges in participant recruitment, enrollment, and

motivation, as highlighted by Brown et al¹⁰⁶ For example, patients having certain medical and psychiatric conditions are generally excluded from the moderate-intensity exercise programs. Moreover, incentives, such as cash, or lack of side effects compared to pharmacotherapy can be used to increase participants' motivation. Secondly, some researchers warn that the advantages of exercise practice are overstated and sport may itself disturb brain function (eg, compromising mental health), especially when intensively performed.^{97,107} Thirdly, Lynch et al¹⁰⁸ and Barha et al¹⁰⁹ reviewed clinical as well as preclinical data suggesting that physical activity efficacy may depend upon the following parameters: (i) individuals (ie, by age and sex), (ii) drug classes (ie, alcohol vs nicotine and cocaine), (iii) stage of the addiction process, and (iv) the exercise conditions tested.

Regarding biological sex, it was argued that women benefited more than men in cognitive outcomes.¹⁰⁹ For example, Colcombe and Kramer¹¹⁰ presented a meta-analysis of 18 RCTs showing that the effect of exercise with an aerobic component on cognition was statistically bigger in samples comprising more than 50% women compared with samples of more than 50% men. Additionally, a 12-month aerobic training resulted in improved attention and memory in older women with MCI, whereas in men, only memory was improved.¹¹¹ Finally, Baker et al¹¹² found that 6 months of high-intensity aerobic exercise had sex bias in cognitive response toward women in the context of mild cognitive impairment. Considering that most of the participants in the studies we reviewed herein were men, it is tempting to speculate that the positive effects of physical exercise would also extend to female METH addicts.

These authors also reviewed METH data, including preclinical studies. For example, Miller et al¹¹³ found that although wheel running (1-hour/day) reduced METH self-administration in rats when it was concurrently available during the acquisition period, it was not effective at decreasing self-administration when it occurred after the acquisition period, indicating that its efficacy may vary with level of exposure or stage of the addiction process. Interestingly, Aarde et al¹¹⁴ demonstrated that 1-day access to a running wheel at the home cage, prior to self-administration sessions, reduced METH intake in rats. Furthermore, Engelmann et al¹¹⁵ showed that although access to a running wheel prior to each self-administration session decreased the acquisition of METH self-administration in rats, a history of unlimited access to a wheel without continued availability enhanced rather than attenuated subsequent acquisition. The impact of physical exercise was also assessed in the withdrawal period in animal models. Recently, Damghani et al¹¹⁶ showed that regular swimming exercise (45 minutes/day, 5 days/week, for 14 days), reduced voluntary METH consumption, and anxiety-like and depression-like behaviors in METH withdrawn rats [rats were exposed to bi-daily doses of METH (2 mg/kg, subcutaneous) over a period of 14 days].

7 | NEUROBIOLOGY OF EXERCISE ON METH ADDICTION MANAGEMENT

The mechanisms of action of physical exercise on the human brain are still poorly known and mostly based on animal studies. Nonetheless, several hypotheses have been proposed to explain the wide variety of beneficial effects attributed to exercise.^{108,117}

One possible neurobiological mechanism mediating the effects of exercise is the modulation of CNS neurochemicals, which play a role in drug addiction and are impaired by METH consumption.^{108,118} In fact, O'Dell et al¹¹⁸ showed that when rats engaged in voluntary aerobic exercise (running wheels) for 3 weeks before and 3 weeks after a binge regimen of METH, exercise significantly ameliorated METH-induced decreases in striatal dopaminergic markers including DAT and TH as well as frontoparietal SERT. This animal dopaminergic data correlate with increased striatal D2/D3 receptor availability in exercised METH abstinence. The apparent correction of striatal dopaminergic deficits operated by physical exercise in METH users in both preclinical and clinical settings seemingly offers better treatment outcomes to METH addicts, namely possible correction of the impulsive temperament which is associated with impaired inhibitory control (please see Section METH toxicodynamics). However, prior exercise in running wheels provided no protection against METH-induced damage to striatal DA terminals in male rats.¹¹⁹ Changes in BDNF levels are also associated to reward-seeking behavior.¹²⁰ Hilburn et al¹²¹ newly found that serum BDNF levels in METH-dependent humans were related to the number of abstinent days since last abuse, but not related to craving and substance use history. More recently, Chen et al¹²² registered constant low BDNF levels in METH abusers, during early withdrawal, when levels of drug seeking are low, when compared to those of the healthy controls. Voluntary aerobic exercise (wheel running) increases BDNF exon IV transcription in rat hippocampus.¹²³ However, METH exposure prior to initiation of aerobic exercise (6 weeks of voluntary wheel running) prevented the increases in both cortical and striatal BDNF seen in saline-treated animals that exercised.¹²⁰ The impact of physical exercise on BDNF levels in METH addicts warrants further evaluation because it was recently demonstrated that patients with serum BDNF levels ≤ 1251.0 pg/mL had higher risk of depression symptoms during METH withdrawal.¹²⁴ We argue that BDNF modifications may contribute to the mood corrections seen in METH addicts that undergo a physical exercise program.

Inoue et al¹²⁵ showed that long-term mild (6-week treadmill running training) rather than intense exercise stimulated adult hippocampal neurogenesis in rats, which was correlated with spatial memory improvement. However, there is limited information regarding the effect of exercise on METH-induced neurogenesis impairment. Recently, Sobieraj et al¹²⁶ provided immunohistochemical analysis of rat brain tissue demonstrating that wheel running during METH withdrawal did not regulate markers of METH neurotoxicity, including neurogenesis, in brain regions involved in relapse to drug seeking. However, reduced METH seeking was associated with running-induced reduction (and normalization) of the number of TH immunoreactive neurons in the periaqueductal gray (PAG). These authors further suggested that wheel running may be preventing certain allostatic changes in the brain reward and stress systems contributing to the negative reinforcement and perpetuation of the addiction cycle. These authors have previously shown that exercise enhanced medial prefrontal cortex gliogenesis (newly generated astrocytes and oligodendrocytes), which was reduced in METH self-administered rats.⁴⁵ On the other

hand, Park et al⁴⁶ demonstrated that METH-induced BBB disruption was concurrent with aberrant adult hippocampal neurogenesis in the DG. Importantly, these authors demonstrated that exercise (wheel running activity) that was introduced post-METH exposure in abstinent mice stabilized the BBB and protected against METH-induced alterations in neurogenesis. These authors highlighted the translational value of their work because exercise is frequently being implemented in dependency treatment of substance-abuse patients.¹²⁷ Further investigations on the potential role of physical exercise in reversing METH-induced gliogenesis, neurogenesis, and BBB impairment are warranted.


Antioxidant effects have also been observed in individuals who perform regular aerobic exercise.¹²⁸ This is consistent with physical exercise ameliorating the reduction-oxidation (redox) balance in the CNS through improvements in the defensive mechanisms of antioxidants.^{129,130} Additionally, Speck et al¹³¹ showed that an adequate frequency of moderate-intensity exercise (treadmill) caused antioxidant changes in the hippocampi of mice. Importantly, voluntary exercise attenuated METH-induced oxidative stress in brain microvessels by raising antioxidant capacity of these capillaries, and protected against BBB disruption, in mice.¹³² Scheme 1 summarizes the proposed neurobiological mechanisms of physical exercise in methamphetamine users.

8 | CONCLUDING REMARKS

To this date, there is no pharmacological treatment directed specifically to METH dependence; thus, it relies on general drugs used in other dependence treatments and on behavioral therapies, with modest outcomes. Physical exercise has been studied as a conjoint therapy in other substance dependences for some time now, with overall positive feedback on sustained abstinence, as seen with tobacco, alcohol, and cannabis use. Only in the past few years, pertinent clinical studies have been conducted addressing specifically METH. METH users that engaged in a physical exercise program exhibited better fitness measures (which were gauged by substantial improvements in aerobic capacity, muscle strength and endurance, body composition, and increased heart rate variability) and showed less depression and anxiety symptoms, lower relapse rates, and sustained abstinence when compared to nonexercised individuals. The neurobiology of physical exercise

in METH users is not fully elucidated and is essentially derived from animal studies. However, positive effects of physical exercise seem to reflect an intricate combination of different players and mechanisms, including neurochemicals, oxidative stress, neurogenesis, gliogenesis, and BBB. Further clinical studies are profoundly needed to confirm reproducibility of previous findings in humans and to dissect the neurobiological basis of physical exercise benefits. We argue that imaging studies (positron emission tomography (PET) and magnetic resonance imaging (MRI) will be instrumental to characterize the impact of PE on oxidative stress (this is a classical METH neurotoxicity hallmark), BBB integrity, and neurogenesis (two recent and exciting hallmarks) in METH addicts.¹³³⁻¹³⁶ Preclinical studies and clinical trials should use physical exercise conjointly with some promising drugs such as mirtazapine, bupropion, and methylphenidate. It is plausible that physical exercise would augment the efficacy of pharmacotherapy. When designing further clinical trials to confirm reproducibility of these beneficial effects of physical exercise on METH addiction, one should attend the following parameters that may help in establishing the most efficient exercise programs: 1—age, 2—sex, 3—neuropsychological consequences of chronic METH use (including cognitive deficits, irritability, agitation, depression, and anxiety symptoms), 4—stage of addiction process (eg, initiation vs relapse), 5—exercise type (anaerobic exercise vs aerobic), duration, and intensity, 5—the need for supervision (structured programs under supervision apparently have better outcomes and most clinical studies happen under residential treatment including psychosocial therapy). As already pinpointed in sections “Biological sex and METH addiction” and “Clinical evidence for physical exercise on METH addiction,” men and women may be affected differently by METH use and by physical exercise. Moreover, it was suggested that physical performance decline similarly in men and women at all ages, and the 1-year age-related declines in performance were about twice as great at 40-year-olds than at 20-year-olds.¹³⁷ Regarding the stages of the addiction process, exercise typically decreases the reinforcing effects of several drugs of abuse, thus preventing drug use initiation, as was shown for alcohol, cigarette, and marijuana use.¹⁰⁸ Therefore, epidemiological studies evaluating the efficacy of physical exercise in METH initiation are warranted. On the other hand, physical exercise may prevent relapse in METH addicts, as suggested by clinical data reviewed herein. The issue of a structured exercise intervention is seemingly relevant. For example, Brown et al¹⁰⁶ outlined two behavioral axes that accompany a moderate-intensity aerobic exercise:

SCHEME 1 Proposed neurobiological mechanisms of physical exercise in methamphetamine users. Physical exercise reverses the neurochemical imbalance, decreases oxidative stress, stabilizes blood-brain barrier, and corrects alterations in neurogenesis and gliogenesis in METH users



(i) group behavioral training component and (ii) an incentive system. In fact, exercise physiologist and psychologist offered to drug addict cognitive and behavioral techniques aiming to increase overall motivation resulting in improved exercise adherence and maintenance. Moreover, participants earned monetary incentives for various levels of adherence to the exercise program. The ultimate goal was to further increase motivation for participation in the physical activity program across the treatment period. For example, Dolezal et al⁹⁸ also referred to a monetary compensation to the METH-dependent individuals in residential treatment during the 12 weeks of study. Regarding exercise type, it has been categorized into aerobic (swimming, running, bicycling) and anaerobic (resistance training, toning) types.¹³⁸ While both types singly improve brain function, the combination of the two types of exercise results in markedly better improvements in cognition than aerobic exercise alone.¹³⁸ Interestingly, clinical data reviewed herein showed that all physical programs employed a combination of aerobic training (jogging and/or walking on a treadmill and cycling) and resistance training. However, Wang et al¹⁰³ used only an aerobic protocol. Nonetheless, it would be interesting to compare the efficacy of both exercise types in the management of METH addicts. Duration and intensity of physical exercise need also to be addressed. In fact, moderate-intensity exercise seems to be associated with more positive effects as highlighted in the Section "Clinical evidence for physical exercise on METH addiction." This is consistent with too much or too little exercise not being advantageous as argued by Alkadhi.¹³⁸ Although the exercise programs reviewed in this manuscript had an 8-week duration, it is reasonable to expect that METH users would retrieve sustained benefits should they continue to exercise beyond the scheduled exercise program. Overall, we propose that exercise-based interventions alone or as a conjoint therapy may be a useful tool for managing METH addiction.

ACKNOWLEDGMENTS

This work was supported by Fundação para a Ciência e Tecnologia (FCT, Portugal) (Strategic Project 2015- UID/NEU/04539/2013), COMPETE-FEDER (POCI-01-0145-FEDER-007400), Centro 2020 Regional Operational Programmes (CENTRO-01-0145-FEDER-000012: HealthyAging2020 and CENTRO-01-0145-FEDER-000008: BrainHealth 2020).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

Frederico Costa Pereira  <http://orcid.org/0000-0002-9381-3320>

REFERENCES

- United Nations Office on Drugs and Crime. *World Drug Report-2016*. New York, NY: United Nations publication; 2016.
- U.S. Department of Justice, Drug Enforcement Administration. 2015 National Drug Threat Assessment Summary. United States: Drug Enforcement Administration; 2015; DEA-DCT-DIR-008-16.
- Courtney KE, Ray LA. Methamphetamine: an update on epidemiology, pharmacology, clinical phenomenology, and treatment literature. *Drug Alcohol Depend*. 2014;143:11-21.
- Warner M, Trinidad JP, Bastian BA, Minimo AM, Hedegaard H. Drugs most frequently involved in drug overdose deaths: United States, 2010-2014. *Natl Vital Stat Rep*. 2016;65:1-15.
- Cho AK, Melega WP. Patterns of methamphetamine abuse and their consequences. *J Addict Dis*. 2002;21:21-34.
- Schifano F, Corkery JM, Cuffolo G. Smokable ("ice", "crystal meth") and non smokable amphetamine-type stimulants: clinical pharmacological and epidemiological issues, with special reference to the UK. *Ann Ist Super Sanita*. 2007;43:110-115.
- Cook CE, Jeffcoat AR, Hill JM, et al. Pharmacokinetics of methamphetamine self-administered to human subjects by smoking S-(+)-methamphetamine hydrochloride. *Drug Metab Dispos*. 1993;21:717-723.
- Rawson RA. Current research on the epidemiology, medical and psychiatric effects, and treatment of methamphetamine use. *J Food Drug Anal*. 2013;21:S77-S81.
- Degenhardt L, Mathers B, Guarinieri M, et al. Methamphetamine use and associated HIV: implications for global policy and public health. *Int J Drug Policy*. 2010;21:347-358.
- Meredith CW, Jaffe C, Ang-Lee K, Saxon AJ. Implications of chronic methamphetamine use: a literature review. *Harv Rev Psychiatry*. 2005;13:141-154.
- Cruickshank CC, Dyer KR. A review of the clinical pharmacology of methamphetamine. *Addiction*. 2009;104:1085-1099.
- Kim I, Oyler JM, Moolchan ET, Cone EJ, Huestis MA. Urinary pharmacokinetics of methamphetamine and its metabolite, amphetamine following controlled oral administration to humans. *Ther Drug Monit*. 2004;26:664-672.
- Homer BD, Solomon TM, Moeller RW, Mascia A, DeRaleau L, Halkitis PN. Methamphetamine abuse and impairment of social functioning: a review of the underlying neurophysiological causes and behavioral implications. *Psychol Bull*. 2008;134:301-310.
- Panenka WJ, Procyshyn RM, Lecomte T, et al. Methamphetamine use: a comprehensive review of molecular, preclinical and clinical findings. *Drug Alcohol Depend*. 2013;129:167-179.
- Rothman RB, Baumann MH. Monoamine transporters and psychostimulant drugs. *Eur J Pharmacol*. 2003;479:23-40.
- Calipari ES, Ferris MJ. Amphetamine mechanisms and actions at the dopamine terminal revisited. *J Neurosci*. 2013;33:8923-8925.
- Han DD, Gu HH. Comparison of the monoamine transporters from human and mouse in their sensitivities to psychostimulant drugs. *BMC Pharmacol*. 2006;6:6.
- Halpin LE, Collins SA, Yamamoto BK. Neurotoxicity of methamphetamine and 3,4-methylenedioxymethamphetamine. *Life Sci*. 2014;97:37-44.
- Dean AC, Groman SM, Morales AM, London ED. An evaluation of the evidence that methamphetamine abuse causes cognitive decline in humans. *Neuropsychopharmacology*. 2013;38:259-274.
- Simon SL, Dean AC, Cordova X, Monterosso JR, London ED. Methamphetamine dependence and neuropsychological functioning: evaluating change during early abstinence. *J Stud Alcohol Drugs*. 2010;71:335-344.
- Volkow ND, Chang L, Wang GJ, et al. Association of dopamine transporter reduction with psychomotor impairment in methamphetamine abusers. *Am J Psychiatry*. 2001;158:377-382.
- Cadet JL, Bisagno V. Neuropsychological consequences of chronic drug use: relevance to treatment approaches. *Front Psychiatry*. 2015;6:189.

23. Thompson PM, Hayashi KM, Simon SL, et al. Structural abnormalities in the brains of human subjects who use methamphetamine. *J Neurosci*. 2004;24:6028-6036.
24. Tobias MC, O'Neill J, Hudkins M, Bartzokis G, Dean AC, London ED. White-matter abnormalities in brain during early abstinence from methamphetamine abuse. *Psychopharmacology*. 2010;209:13-24.
25. Nestor LJ, Ghahremani DG, Monterosso J, London ED. Prefrontal hypoactivation during cognitive control in early abstinent methamphetamine-dependent subjects. *Psychiatry Res*. 2011;194:287-295.
26. Kohno M, Morales AM, Ghahremani DG, Hellemann G, London ED. Risky decision making, prefrontal cortex, and mesocorticolimbic functional connectivity in methamphetamine dependence. *JAMA Psychiatry*. 2014;71:812-820.
27. London ED, Simon SL, Berman SM, et al. Mood disturbances and regional cerebral metabolic abnormalities in recently abstinent methamphetamine abusers. *Arch Gen Psychiatry*. 2004;61:73-84.
28. Payer DE, Lieberman MD, Monterosso JR, Xu J, Fong TW, London ED. Differences in cortical activity between methamphetamine-dependent and healthy individuals performing a facial affect matching task. *Drug Alcohol Depend*. 2008;93:93-102.
29. London ED, Kohno M, Morales AM, Ballard ME. Chronic methamphetamine abuse and corticostriatal deficits revealed by neuroimaging. *Brain Res*. 2015;1628:174-185.
30. McCann UD, Wong DF, Yokoi F, Villemagne V, Dannals RF, Ricaurte GA. Reduced striatal dopamine transporter density in abstinent methamphetamine and methcathinone users: evidence from positron emission tomography studies with [¹¹C]WIN-35,428. *J Neurosci*. 1998;18:8417-8422.
31. McCann UD, Kuwabara H, Kumar A, et al. Persistent cognitive and dopamine transporter deficits in abstinent methamphetamine users. *Synapse*. 2008;62:91-100.
32. Sekine Y, Minabe Y, Ouchi Y, et al. Association of dopamine transporter loss in the orbitofrontal and dorsolateral prefrontal cortices with methamphetamine-related psychiatric symptoms. *Am J Psychiatry*. 2003;160:1699-1701.
33. Lee B, London ED, Poldrack RA, et al. Striatal dopamine D2/D3 receptor availability is reduced in methamphetamine dependence and is linked to impulsivity. *J Neurosci*. 2009;29:14734-14740.
34. Zorick T, Lee B, Mandelkern MA, et al. Low striatal dopamine receptor availability linked to caloric intake during abstinence from chronic methamphetamine abuse. *Mol Psychiatry*. 2011;17:569-571.
35. Kitamura O, Tokunaga I, Gotohda T, Kubo S. Immunohistochemical investigation of dopaminergic terminal markers and caspase-3 activation in the striatum of human methamphetamine users. *Int J Legal Med*. 2007;121:163-168.
36. Moszczynska A, Fitzmaurice P, Ang L, et al. Why is parkinsonism not a feature of human methamphetamine users? *Brain*. 2004;127:363-370.
37. Wilson JM, Kalasinsky KS, Levey AI, et al. Striatal dopamine nerve terminal markers in human, chronic methamphetamine users. *Nat Med*. 1996;2:699-703.
38. Kish SJ, Fitzmaurice PS, Boileau I, et al. Brain serotonin transporter in human methamphetamine users. *Psychopharmacology*. 2009;202:649-661.
39. Solhi H, Malekiran A, Kazemifar AM, Sharifi F. Oxidative stress and lipid peroxidation in prolonged users of methamphetamine. *Drug Metab Lett*. 2014;7:79-82.
40. Northrop NA, Yamamoto BK. Persistent neuroinflammatory effects of serial exposure to stress and methamphetamine on the blood-brain barrier. *J Neuroimmune Pharmacol*. 2012;7:951-968.
41. Park M, Hennig B, Toborek M. Methamphetamine alters occludin expression via NADPH oxidase-induced oxidative insult and intact caveolae. *J Cell Mol Med*. 2012;16:362-375.
42. Ramirez SH, Potula R, Fan S, et al. Methamphetamine disrupts blood-brain barrier function by induction of oxidative stress in brain endothelial cells. *J Cereb Blood Flow Metab*. 2009;29:1933-1945.
43. Sharma HS, Kiyatkin EA. Rapid morphological brain abnormalities during acute methamphetamine intoxication in the rat: an experimental study using light and electron microscopy. *J Chem Neuroanat*. 2009;37:18-32.
44. Ago M, Ago K, Hara K, Kashimura S, Ogata M. Toxicological and histopathological analysis of a patient who died nine days after a single intravenous dose of methamphetamine: a case report. *Leg Med*. 2006;8:235-239.
45. Mandyam CD, Wee S, Eisch AJ, Richardson HN, Koob GF. Methamphetamine self-administration and voluntary exercise have opposing effects on medial prefrontal cortex gliogenesis. *J Neurosci*. 2007;27:11442-11450.
46. Park M, Levine H, Toborek M. Exercise protects against methamphetamine-induced aberrant neurogenesis. *Sci Rep*. 2016;6:34111.
47. Recinto P, Samant AR, Chavez G, et al. Levels of neural progenitors in the hippocampus predict memory impairment and relapse to drug seeking as a function of excessive methamphetamine self-administration. *Neuropsychopharmacology*. 2012;37:1275-1287.
48. Galinato MH, Lockner JW, Fannon-Pavlich MJ, et al. A synthetic small-molecule Isoxazole-9 protects against methamphetamine relapse. *Mol Psychiatry*. 2017;20:4226-4235.
49. Mooney LJ, Glasner-Edwards S, Rawson RA, Ling W. Medical effects of methamphetamine use. In: Roll JM, Rawson RA, Ling W, Shoptaw S, eds. *Methamphetamine Addiction: From Basic Science to Treatment*. New York, NY: Guilford Press; 2009:117-142.
50. Moszczynska A, Callan SP. Molecular, behavioral, and physiological consequences of methamphetamine neurotoxicity: implications for Treatment. *J Pharmacol Exp Ther*. 2017;362:474-488.
51. Richards JR, Bretz SW, Johnson EB, Turnipseed SD, Brofeldt BT, Derlet RW. Methamphetamine abuse and emergency department utilization. *West J Med*. 1999a;170:198-202.
52. Richards JR, Johnson EB, Stark RW, Derlet RW. Methamphetamine abuse and rhabdomyolysis in the ED: a 5-year study. *Am J Emerg Med*. 1999b;17:681-685.
53. Mendelson J, Uemura N, Harris D, et al. Human pharmacology of the methamphetamine stereoisomers. *Clin Pharmacol Ther*. 2006;80:403-420.
54. Jones ES, Rayner BL. Hypertension, end-stage renal disease and mesangiocapillary glomerulonephritis in methamphetamine users. *S Afr Med J*. 2015;105:199-201.
55. Bowyer JF, Hanig JP. Amphetamine- and methamphetamine-induced hyperthermia: implications of the effects produced in brain vasculature and peripheral organs to forebrain neurotoxicity. *Temperature*. 2014;1:172-182.
56. Cantwell B, McBride AJ. Self detoxication by amphetamine dependent patients: a pilot study. *Drug Alcohol Depend*. 1998;49:157-163.
57. Hser YI, Evans E, Huang YC. Treatment outcomes among women and men methamphetamine abusers in California. *J Subst Abuse Treat*. 2005;28:77-85.
58. Rawson RA, Marinelli-Casey P, Anglin MD, et al. A multi-site comparison of psychosocial approaches for the treatment of methamphetamine dependence. *Addiction*. 2004;99:708-717.
59. Smith K. *The CBHSQ Report*. Rockville, MD: Substance Abuse and Mental Health Services Administration (US); 2013-2014.
60. Cohen JB, Greenberg R, Uri J, Halpin M, Zweben JE. Women with methamphetamine dependence: research on etiology and treatment. *J Psychoactive Drugs*. 2007;39(Suppl 4):347-351.
61. Simpson JL, Grant KM, Daly PM, Kelley SG, Carlo G, Bevins RA. Psychological burden and gender differences in

- methamphetamine-dependent individuals in treatment. *J Psychoactive Drugs*. 2016;48:261-269.
62. Kogachi S, Chang L, Alicata D, Cunningham E, Ernst T. Sex differences in impulsivity and brain morphometry in methamphetamine users. *Brain Struct Funct*. 2017;222:215-227.
 63. Rawson RA, Huber A, Brethen P, et al. Status of methamphetamine users 2-5 years after outpatient treatment. *J Addict Dis*. 2002;21:107-119.
 64. Brecht ML, Lovinger K, Herbeck DM, Urada D. Patterns of treatment utilization and methamphetamine use during first 10 years after methamphetamine initiation. *J Subst Abuse Treat*. 2013;44:548-556.
 65. Downey LA, Loftis JM. Altered energy production, lowered antioxidant potential, and inflammatory processes mediate CNS damage associated with abuse of the psychostimulants MDMA and methamphetamine. *Eur J Pharmacol*. 2014;727:125-129.
 66. Brecht ML, Herbeck D. Time to relapse following treatment for methamphetamine use: a long-term perspective on patterns and predictors. *Drug Alcohol Depend*. 2014;139:18-25.
 67. Ling W, Rawson R, Shoptaw S, Ling W. Management of methamphetamine abuse and dependence. *Curr Psychiatry Rep*. 2006;8:345-354.
 68. Ciccarone D. Stimulant abuse: pharmacology, cocaine, methamphetamine, treatment, attempts at pharmacotherapy. *Prim Care*. 2011;38:41-58.
 69. Brensilver M, Heinzerling KG, Shoptaw S. Pharmacotherapy of amphetamine-type stimulant dependence: an update. *Drug Alcohol Rev*. 2013;32:449-460.
 70. Karila L, Weinstein A, Aubin HJ, Benyamina A, Reynaud M, Batki SL. Pharmacological approaches to methamphetamine dependence: a focused review. *Br J Clin Pharmacol*. 2010;69:578-592.
 71. Worley MJ, Heinzerling KG, Roche DJ, Shoptaw S. Ibudilast attenuates subjective effects of methamphetamine in a placebo-controlled inpatient study. *Drug Alcohol Depend*. 2016;162:245-250.
 72. Zanetidou S, Belvederi Murri M, Menchetti M, et al. Physical exercise for late-life depression: customizing an intervention for primary care. *J Am Geriatr Soc*. 2017;65:348-355.
 73. Colfax GN, Santos GM, Das M, et al. Mirtazapine to reduce methamphetamine use: a randomized controlled trial. *Arch Gen Psychiatry*. 2011;68:1168-1175.
 74. De La Garza R 2nd, Newton TF, Haile CN, et al. Rivastigmine reduces "likely to use methamphetamine" in methamphetamine-dependent volunteers. *Prog Neuropsychopharmacol Biol Psychiatry* 2012;37:141-146.
 75. Elkashef AM, Rawson RA, Anderson AL, et al. Bupropion for the treatment of methamphetamine dependence. *Neuropsychopharmacology*. 2008;33:1162-1170.
 76. Elkashef A, Kahn R, Yu E, et al. Topiramate for the treatment of methamphetamine addiction: a multi-center placebo-controlled trial. *Addiction*. 2012;107:1297-1306.
 77. Galloway GP, Buscemi R, Coyle JR, et al. A randomized, placebo-controlled trial of sustained-release dextroamphetamine for treatment of methamphetamine addiction. *Clin Pharmacol Ther*. 2011;89:276-282.
 78. Grant JE, Odlaug BL, Kim SW. A double-blind, placebo-controlled study of n-acetyl cysteine plus naltrexone for methamphetamine dependence. *Eur Neuropsychopharmacol*. 2010;20:823-828.
 79. Heinzerling KG, Shoptaw S, Peck JA, et al. Randomized, placebo-controlled trial of baclofen and gabapentin for the treatment of methamphetamine dependence. *Drug Alcohol Depend*. 2006;85:177-184.
 80. Heinzerling KG, Swanson AN, Kim S, et al. Randomized, double-blind, placebo-controlled trial of modafinil for the treatment of methamphetamine dependence. *Drug Alcohol Depend*. 2010;109:20-29.
 81. Morley KC, Cornish JL, Faingold A, Wood K, Haber PS. Pharmacotherapeutic agents in the treatment of methamphetamine dependence. *Expert Opin Investig Drugs*. 2017;26:563-578.
 82. Mousavi SG, Sharbafchi MR, Salehi M, Peykanpour M, Karimian Sichani N, Maracy M. The efficacy of n-acetylcysteine in the treatment of methamphetamine dependence: a double-blind controlled, crossover study. *Arch Iran Med*. 2015;18:28-33.
 83. Rezaei F, Emami M, Zahed S, Morabbi MJ, Farahzadi M, Akhondzadeh S. Sustained-release methylphenidate in methamphetamine dependence treatment: a double-blind and placebo-controlled trial. *Daru*. 2015;23:2.
 84. Sulaiman AH, Gill JS, Said MA, Zainal NZ, Hussein HM, Guan NC. A randomized, placebo-controlled trial of aripiprazole for the treatment of methamphetamine dependence and associated psychosis. *Int J Psychiatry Clin Pract*. 2013;17:131-138.
 85. Verrico CD, Mahoney JJ 3rd, Thompson-Lake DG, Bennett RS, Newton TF, De La Garza R 2nd. Safety and efficacy of varenicline to reduce positive subjective effects produced by methamphetamine in methamphetamine-dependent volunteers. *Int J Neuropsychopharmacol*. 2014;17:223-233.
 86. Verrico CD, Haile CN, De La Garza R 2nd, Grasing K, Kosten TR, Newton TF. Subjective and cardiovascular effects of intravenous methamphetamine during perindopril maintenance: a randomized, double-blind, placebo-controlled human laboratory study. *Int J Neuropsychopharmacol* 2016;19:pyw029.
 87. Lee NK, Rawson RA. A systematic review of cognitive and behavioural therapies for methamphetamine dependence. *Drug Alcohol Rev*. 2008;27:309-317.
 88. Rawson RA, Shoptaw SJ, Obert JL, et al. An intensive outpatient approach for cocaine abuse treatment. The Matrix model. *J Subst Abuse Treat*. 1995;12:117-127.
 89. Marinelli-Casey P, Gonzales R, Hillhouse M, et al. Drug court treatment for methamphetamine dependence: treatment response and posttreatment outcomes. *J Subst Abuse Treat*. 2008;34:242-248.
 90. Shearer J. Psychosocial approaches to psychostimulant dependence: a systematic review. *Psychosocial approaches to psychostimulant dependence: a systematic review. J Subst Abuse Treat*. 2007;32:41-52.
 91. Baicy K, London ED. Corticolimbic dysregulation and chronic methamphetamine abuse. *Addiction*. 2007;102(Suppl 1):5-15.
 92. Mooney LJ, Cooper C, London ED, et al. Exercise for methamphetamine dependence: rationale, design, and methodology. *Contemp Clin Trials*. 2014;37:139-147.
 93. Rawson RA, Chudzynski J, Gonzales R, et al. The impact of exercise on depression and anxiety symptoms among abstinent methamphetamine-dependent individuals in a residential treatment setting. *J Subst Abuse Treat*. 2015a;57:36-40.
 94. Pekmezi DW, Carr LJ, Barbera B, Marcus BH. The role of physical activity in treatment of substance use disorders. In: Meyer AL, Gullotta TP, eds. *Physical Activity Across the Lifespan*. New York, NY: Springer; 2012:171-191.
 95. Linke SE, Ussher M. Exercise-based treatments for substance use disorders: evidence, theory, and practicality. *Am J Drug Alcohol Abuse*. 2015;41:7-15.
 96. Buchowski MS, Meade NN, Charboneau E, et al. Aerobic exercise training reduces cannabis craving and use in non-treatment seeking cannabis-dependent adults. *PLoS One*. 2011;6:e17465.
 97. Neale J, Nettleton S, Pickering L. Heroin users' views and experiences of physical activity, sport and exercise. *Int J Drug Policy*. 2012;23:120-127.
 98. Dolezal BA, Chudzynski J, Storer TW, et al. Eight weeks of exercise training improves fitness measures in methamphetamine-dependent individuals in residential treatment. *J Addict Med*. 2013;7:122-128.
 99. Dolezal BA, Chudzynski J, Dickerson D, et al. Exercise training improves heart rate variability after methamphetamine dependency. *Med Sci Sports Exerc*. 2014;46:1057-1066.
 100. Robertson CL, Ishibashi K, Chudzynski J, et al. Effect of exercise training on striatal dopamine D2/D3 receptors in methamphetamine users during behavioral treatment. *Neuropsychopharmacology*. 2016;41:1629-1636.
 101. Rawson RA, Chudzynski J, Mooney L, et al. Impact of an exercise intervention on methamphetamine use outcomes post-residential treatment care. *Drug Alcohol Depend*. 2015b;156:21-28.

102. Haglund M, Ang A, Mooney L, et al. Predictors of depression outcomes among abstinent methamphetamine-dependent individuals exposed to an exercise intervention. *Am J Addict*. 2015;24:246-251.
103. Wang D, Zhou C, Zhao M, Wu X, Chang YK. Dose-response relationships between exercise intensity, cravings, and inhibitory control in methamphetamine dependence: an ERPs study. *Drug Alcohol Depend*. 2016;161:331-339.
104. Wang D, Wang C, Zhou YK, Chang YK. Acute exercise ameliorates craving and inhibitory deficits in methamphetamine: an ERP study. *Physiol Behav*. 2015;147:38-46.
105. Wang D, Zhu T, Zhou C, Chang YK. Aerobic exercise training ameliorates craving and inhibitory control in methamphetamine dependencies: a randomized controlled trial and event-related potential study. *Psychol Sport Exerc*. 2017;30:82-90.
106. Brown RA, Abrantes AM, Read JP, et al. A Pilot study of aerobic exercise as an adjunctive treatment for drug dependence. *Ment Health Phys Act*. 2010;3:27-34.
107. Armstrong LE, VanHeest JL. The unknown mechanism of the overtraining syndrome: clues from depression and psychoneuroimmunology. *Sports Med*. 2002;32:185-209.
108. Lynch WJ, Peterson AB, Sanchez V, Abel J, Smith MA. Exercise as a novel treatment for drug addiction: a neurobiological and stage-dependent hypothesis. *Neurosci Biobehav Rev*. 2013;37:1622-1644.
109. Barha CK, Davis JC, Falck RS, Nagamatsu LS, Liu-Ambrose T. Sex differences in exercise efficacy to improve cognition: a systematic review and meta-analysis of randomized controlled trials in older humans. *Front Neuroendocrinol*. 2017;46:71-85.
110. Colcombe S, Kramer AF. Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol Sci*. 2003;14:125-130.
111. van Uffelen JG, Chinapaw MJ, van Mechelen W, Hopman-Rock M. Walking or vitamin B for cognition in older adults with mild cognitive impairment? A randomised controlled trial. *Br J Sports Med*. 2008;42:344-351.
112. Baker LD, Frank LL, Foster-Schubert K, et al. Effects of aerobic exercise on mild cognitive impairment: a controlled trial. *Arch Neurol*. 2010;67:71-79.
113. Miller ML, Vaillancourt BD, Wright MJ Jr, et al. Reciprocal inhibitory effects of intravenous d-methamphetamine self-administration and wheel activity in rats. *Drug Alcohol Depend*. 2012;121:90-96.
114. Aarde SM, Miller ML, Creehan KM, Vandewater SA, Taffe MA. One day access to a running wheel reduces self-administration of D-methamphetamine, MDMA and methylone. *Drug Alcohol Depend*. 2015;151:151-158.
115. Engelmann AJ, Aparicio MB, Kim A, et al. Chronic wheel running reduces maladaptive patterns of methamphetamine intake: regulation by attenuation of methamphetamine-induced neuronal nitric oxide synthase. *Brain Struct Funct*. 2014;219:657-672.
116. Damghani F, Bigdeli I, Miladi-Gorji H, Fadaei A. Swimming exercise attenuates psychological dependence and voluntary methamphetamine consumption in methamphetamine withdrawn rats. *Iran J Basic Med Sci*. 2016;19:594-600.
117. Fontes-Ribeiro CA, Marques E, Pereira FC, Silva AP, Macedo TR. May exercise prevent addiction? *Curr Neuropharmacol*. 2011;9:45-48.
118. O'Dell SJ, Galvez BA, Ball AJ, Marshall JF. Running wheel exercise ameliorates methamphetamine-induced damage to dopamine and serotonin terminals. *Synapse*. 2012;66:71-80.
119. O'Dell SJ, Marshall JF. Running wheel exercise before a binge regimen of methamphetamine does not protect against striatal dopaminergic damage. *Synapse*. 2014;68:419-425.
120. Thompson AB, Stolyarova A, Ying Z, Zhuang Y, Gómez-Pinilla F, Izquierdo A. Methamphetamine blocks exercise effects on Bdnf and Drd2 gene expression in frontal cortex and striatum. *Neuropharmacology*. 2015;99:658-664.
121. Hilburn C, Nejtek VA, Underwood WA, et al. Is serum brain-derived neurotrophic factor related to craving for or use of alcohol, cocaine, or methamphetamine? *Neuropsychiatr Dis Treat*. 2011;7:357-364.
122. Chen PH, Huang MC, Lai YC, Chen PY, Liu HC. Serum brain-derived neurotrophic factor levels were reduced during methamphetamine early withdrawal. *Addict Biol*. 2014;19:482-485.
123. Gomez-Pinilla F, Zhuang Y, Feng J, Ying Z, Fan G. Exercise impacts brain-derived neurotrophic factor plasticity by engaging mechanisms of epigenetic regulation. *Eur J Neurosci*. 2011;33:383-390.
124. Ren W, Luan X, Zhang J, et al. Brain-derived neurotrophic factor levels and depression during methamphetamine withdrawal. *Affect Disord*. 2017;221:165-171.
125. Inoue K, Okamoto M, Shibato J, et al. Long-term mild, rather than intense, exercise enhances adult hippocampal neurogenesis and greatly changes the transcriptomic profile of the hippocampus. *PLoS One*. 2015;10:e0128720.
126. Sobieraj JC, Kim A, Fannon MJ, Mandyam CD. Chronic wheel running-induced reduction of extinction and reinstatement of methamphetamine seeking in methamphetamine dependent rats is associated with reduced number of periaqueductal gray dopamine neurons. *Brain Struct Funct*. 2016;221:261-276.
127. Wang D, Wang Y, Wang Y, Li R, Zhou C. Impact of physical exercise on substance use disorders: a meta-analysis. *PLoS One*. 2014;9:e110728.
128. Radak Z, Chung HY, Goto S. Systemic adaptation to oxidative challenge induced by regular exercise. *Free Radic Biol Med*. 2008;44:153-159.
129. Aguiar AS Jr, Tuon T, Soares FS, da Rocha LG, Silveira PC, Pinho RA. The effect of n-acetylcysteine and deferoxamine on exercise-induced oxidative damage in striatum and hippocampus of mice. *Neurochem Res*. 2008;33:729-736.
130. Camiletti-Moiron D, Aparicio VA, Aranda P, Radak Z. Does exercise reduce brain oxidative stress? A systematic review. *Scand J Med Sci Sports*. 2013;23:e202-e212.
131. Speck AE, Tromm CB, Pozzi BG, et al. The dose-dependent antioxidant effects of physical exercise in the hippocampus of mice. *Neurochem Res*. 2014;39:1496-1501.
132. Toborek M, Seelbach MJ, Rashid CS, et al. Voluntary exercise protects against methamphetamine-induced oxidative stress in brain microvasculature and disruption of the blood-brain barrier. *Mol Neurodegener*. 2013;8:22.
133. Tamura Y, Kataoka Y. PET imaging of neurogenic activity in the adult brain: toward in vivo imaging of human neurogenesis. *Neurogenesis*. 2017;4:e1281861.
134. Neishi H, Ikawa M, Okazawa H, et al. Precise evaluation of striatal oxidative stress corrected for severity of dopaminergic neuronal degeneration in patients with Parkinson's disease: a study with 62Cu-ATSM PET and 123I-FP-CIT SPECT. *Eur Neurol*. 2017;78:161-168.
135. Pollak TA, Drndarski S, Stone JM, David AS, McGuire P, Abbott NJ. The blood-brain barrier in psychosis. *Lancet Psychiatry*. 2017; [Epub ahead of print]. [https://doi.org/10.1016/s2215-0366\(17\)30293-6](https://doi.org/10.1016/s2215-0366(17)30293-6).
136. Veksler R, Shelef I, Friedman A. Blood-brain barrier imaging in human neuropathologies. *Arch Med Res*. 2014;45:646-652.
137. Bongard V, McDermott AY, Dallal GE, Schaefer EJ. Effects of age and gender on physical performance. *Age*. 2007;29:77-85.
138. Alkadhi KA. Exercise as a positive modulator of brain function. *Mol Neurobiol*. 2017;6(Suppl 3):765-783.

How to cite this article: Morais APD, Pita IR, Fontes-Ribeiro CA, Pereira FC. The neurobiological mechanisms of physical exercise in methamphetamine addiction. *CNS Neurosci Ther*. 2017;00:1-13. <https://doi.org/10.1111/cns.12788>